

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2-Amino-N-(2-methoxyphenyl)-4,5-dimethylthiophene-3-carboxamide

K. Chandra Kumar,<sup>a</sup> M. K. Kokila,<sup>b\*</sup> Puttaraja,<sup>b</sup>  
J. Saravanan<sup>c</sup> and Manohar V. Kulkarni<sup>d</sup><sup>a</sup>Department of Engineering Physics, HKBK College of Engineering, Nagawara, Bangalore 560 045, Karnataka, India, <sup>b</sup>Department of Physics, Bangalore University, Bangalore 560 056, Karnataka, India, <sup>c</sup>PES College of Pharmacy, Hanumanthanagar, Bangalore 560 050, Karnataka, India, and <sup>d</sup>Department of Chemistry, Karnatak University, Dharwad 580 003, Karnataka, India  
Correspondence e-mail: [prmkkgroup@gmail.com](mailto:prmkkgroup@gmail.com)


Received 5 June 2008; accepted 16 June 2008

Key indicators: single-crystal X-ray study;  $T = 291$  K; mean  $\sigma(\text{C}—\text{C}) = 0.005$  Å;  $R$  factor = 0.054;  $wR$  factor = 0.143; data-to-parameter ratio = 14.4.

In the title compound,  $\text{C}_{14}\text{H}_{16}\text{N}_2\text{O}_2\text{S}$ , the two aromatic rings make a dihedral angle of  $13.9(1)^\circ$ . The crystal structure is stabilized by both inter- and intramolecular  $\text{N}—\text{H}\cdots\text{O}$ ,  $\text{C}—\text{H}\cdots\text{O}$  and  $\text{C}—\text{H}\cdots\text{N}$  hydrogen bonds.

## Related literature

For related literature, see: Gewald *et al.* (1966); Cohen *et al.* (1977); Csaszar & Morvay (1983); Lakshmi *et al.* (1985); Mohan & Saravanan (2003); Bruns *et al.* (1990).



## Experimental

## Crystal data

 $\text{C}_{14}\text{H}_{16}\text{N}_2\text{O}_2\text{S}$   
 $M_r = 276.35$   
Monoclinic,  $P2_1/n$   
 $a = 8.606(2)$  Å  
 $b = 7.5193(19)$  Å  
 $c = 21.297(5)$  Å  
 $\beta = 100.599(5)^\circ$  $V = 1354.7(6)$  Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.24$  mm<sup>-1</sup>  
 $T = 291(2)$  K  
 $0.45 \times 0.35 \times 0.28$  mm

## Data collection

Bruker SMART CCD area-detector  
diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.908$ ,  $T_{\max} = 0.937$ 9834 measured reflections  
2514 independent reflections  
1503 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.051$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.053$   
 $wR(F^2) = 0.142$   
 $S = 0.99$   
2514 reflections175 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.25$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.18$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
$\text{N1}—\text{H1A}\cdots\text{O1}$	0.86	2.15	2.724 (3)	124
$\text{N1}—\text{H1B}\cdots\text{O1}^i$	0.86	2.23	3.009 (4)	151
$\text{N2}—\text{H2}\cdots\text{O2}$	0.86	2.15	2.565 (3)	109
$\text{C8}—\text{H8}\cdots\text{O1}$	0.93	2.30	2.874 (4)	119

Symmetry code: (i)  $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$ .

Data collection: *SMART* (Bruker, 1998); cell refinement: *SMART*; data reduction: *SAINT* (Bruker, 1998); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *PARST* (Nardelli, 1995) and *PLATON* (Spek, 2003).

The authors are grateful to Professor T. N. Guru Row, Indian Institute of Science, and the Department of Science and Technology, India, for the data collection using the CCD facility, and Bangalore University. CK thanks the Management, Administrator and Principal of HKBK College of Engineering for encouragement and support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2721).

## References

- Bruker (1998). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruns, R. F., Fergus, J. H., Coughenour, L. L., Courtland, G. E., Pugsley, T. A., Dodd, J. H. & Tinney, F. J. (1990). *Mol. Pharmacol.* **38**, 950–958.
- Cohen, V. I., Rist, N. & Duponchel, C. (1977). *J. Pharm. Sci.* **66**, 1332–1334.
- Csaszar, J. & Morvay, J. (1983). *Acta Pharm. Hung.* **53**, 121–128.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Gewald, K., Schinke, E. & Botcher, H. (1966). *Chem. Ber.* **99**, 94–100.
- Lakshmi, V. V., Sridhar, P. & Polsa, H. (1985). *Indian J. Pharm. Sci.* **23**, 327–336.
- Mohan, S. & Saravanan, J. (2003). *Asian J. Chem.* **15**, 67–70.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

**supplementary materials**



## Figures




Fig. 1. The molecular structure of (I), with atom labels and 50% probability displacement ellipsoids for non-H atoms. Dashed lines indicate intramolecular hydrogen bonds; H atoms not involved in hydrogen bonding have been omitted.




Fig. 2. The packing of (I), viewed down the *a* axis shows molecules connected by N—H...O hydrogen bonds (dashed lines). H atoms not involved in hydrogen bonding have been omitted.

## 2-Amino-N-(2-methoxyphenyl)-4,5-dimethylthiophene-3-carboxamide

### Crystal data

$C_{14}H_{16}N_2O_2S$

$M_r = 276.35$

Monoclinic,  $P2_1/n$

Hall symbol:  $-P2_1/n$

$a = 8.606$  (2) Å

$b = 7.5193$  (19) Å

$c = 21.297$  (5) Å

$\beta = 100.599$  (5)°

$V = 1354.7$  (6) Å<sup>3</sup>

$Z = 4$

$F_{000} = 584$

$D_x = 1.355$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation

$\lambda = 0.71073$  Å

Cell parameters from 670 reflections

$\theta = 2.0$ – $28.5^\circ$

$\mu = 0.24$  mm<sup>-1</sup>

$T = 291$  (2) K

Block, yellow

$0.45 \times 0.35 \times 0.28$  mm













